

On area and entropy of a black hole

A. Alekseev[†], A. P. Polychronakos^{†*}, M. Smedbäck[†]

[†] *Institute for Theoretical Physics, Uppsala University, Box 803, S-75108, Uppsala, Sweden*

^{*} *Theoretical Physics Dept., University of Ioannina, 45110 Ioannina, Greece*

e-mails: alekseev@teorfys.uu.se, poly@teorfys.uu.se, mikael@teorfys.uu.se

(April 2000)

We consider a model of a black hole consisting of a number of elementary components. Examples of such models occur in the Ashtekar's approach to canonical Quantum Gravity and in M-theory. We show that treating the elementary components as completely distinguishable leads to the area law for the black hole entropy. Contrary to previous results, we show that no Bose condensation occurs, the area has big local fluctuations and that in the framework of canonical Quantum Gravity the area of the black hole horizon is equidistantly quantized.

PACS-98: 04.70.Dy, 11.15.-q

In the framework of Quantum Gravity black holes are treated as quantum objects. As such, they are characterized by quantum numbers: mass, electric charge, angular momentum *etc.* For Schwarzschild black holes (neutral, nonrotating) the only quantum number which is left is the mass M . It is related to the area A of the black hole horizon by formula, $A = 16\pi G^2 M^2 / c^4$, where G is the gravitational constant. Important questions in black hole physics are what the spectrum of A looks like and what the degeneracies of states are for a given value of A .

In the absence of a definitive Quantum Gravity theory, the answers to these questions depend on the model of a black hole. In several approaches one assumes that a black hole consists of elementary components contributing additively to its area. For instance, in the Ashtekar's approach (see *e.g.* [1,2]) these are Wilson lines of the Ashtekar's connection A_μ^a . In the M -theory approach [3] these are D₀-branes. Each of the elementary components can be in a number of states. An elementary component in the state n gives a contribution A_n to the total area, and has a degeneracy $g(n)$. The eigenvalues of the area operator then acquire the form $A = \sum_{n=1}^{\infty} N_n A_n$, where N_n is the number of elementary components in the state n . The multiplicity of states with area A is $\Omega(A)$ and its asymptotics when A is macroscopically large defines the entropy of the black hole, $S(A) = k \ln \Omega(A)$.

The spectrum of the area A and the behaviour of the entropy $S(A)$ depend on the spectrum of the elementary area A_n , the elementary multiplicities $g(n)$ and the degree of distinguishability of the components. We will discuss the form of A_n in the framework of canonical Quantum Gravity [1,2]. The classical result in this field is that the index n can be associated with a spin j of an $SU(2)$ representation. Then, j takes integer and half-integer values and according to [5] $A_j = \sqrt{j(j+1)}$ in some units. We argue that instead

$$A_j = j + \frac{1}{2}. \quad (1)$$

In particular, this implies that the spectrum of the area operator is equidistant. Such a situation was first considered by Bekenstein and Mukhanov [6]. Their analysis shows that such a spectrum implies certain specific properties of the black hole radiation: there exists an energy quantum, $\hbar\omega_0$, and the energy can be radiated only in integer multiples of this energy quantum.

Furthermore, we will consider the issue of black hole entropy. It will be seen that the *area law* $S \propto A$ is to a large extent generic. A more subtle issue is what the statistically preferable state of the system is. The analysis of [7] in the case of Ashtekar gravity and of [8] in the case of black holes composed of D₀-branes showed that the most probable configuration has all the elementary components in the same multiplet. This can be viewed as Bose condensation. We show instead that for completely distinguishable elementary components one obtains a Gibbs distribution for the N_n 's with many of them nonvanishing in the most probable configuration. Our result implies that the area operator has strong fluctuations, $\Delta A \sim \sqrt{A}$, whereas in the case of Bose condensation the fluctuation of area would have been strongly suppressed.

Next we turn to the issue of the area spectrum in canonical Quantum Gravity. One of the virtues of Ashtekar's formulation is that instead of the metric $g_{\mu\nu}$ one deals with the gauge field A_μ^a [1,2]. In the Euclidean gravity the gauge group is $SU(2)$ and the isotopic index a takes values 1, 2, 3 corresponding to three Pauli matrices. In the Hamiltonian formulation one uses the spatial components A_i^a as generalized coordinates and the components of the electric field, $E_i = \partial_0 A_i - \partial_i A_0 - [A_0, A_i]$ as conjugate momenta. It is convenient to view the gauge

field as a 1-form $A = A_i dx^i$, and the conjugate momenta E_i as components of a 2-form, $E = \frac{1}{2} \varepsilon_{ijk} E_i dx^j dx^k$.

The physical Hilbert space is obtained by imposing the Gauss law constraint, the diffeomorphism constraint and the Hamiltonian constraint (the Wheeler-DeWitt equation). While constructing wave functionals which annihilate all those constraints proves to be a difficult task [1], one believes that the spectrum of the area operator is already captured by studying the ‘kinematical Hilbert space’ containing the Wilson lines [9],

$$W_\Gamma^j(A) = \text{Tr} P \exp \left(\int_\Gamma A^a T_j^a \right), \quad (2)$$

where Γ is a closed contour, j is a positive integer or half-integer and T_j^a are generators of the Lie algebra $su(2)$ in the representation with spin j .

In terms of the Ashtekar’s variables, the area operator corresponding to the 2-dimensional surface Σ acquires the form [5],

$$\mathcal{A}_\Sigma = \int_\Sigma d^2x \sqrt{\text{Tr} E^2}, \quad (3)$$

where the integrand is the natural density which can be integrated over the 2-dimensional surface Σ . The Wilson lines W_Γ^j are eigenstates of the operators $\mathcal{A}_\Sigma$, at least when all the intersections of Γ and Σ are transversal. Moreover, each intersection gives a contribution A_j which depends only on the spin j of the Wilson line. It is our next task to compute the numbers A_j .

Canonical quantization suggests that in the A -representation the conjugate momentum E_i^a act as derivatives, $E_i^a(x) = -i\delta/\delta A_i^a(x)$. Ignoring the singularity arising from the coincident arguments in the expression $\sqrt{E_i^a(x)E_i^a(x)}$, one easily obtains

$$\mathcal{A}_\Sigma W_\Gamma(A) = \sum_p \text{Tr} \sqrt{T_j^a T_j^a} P \exp \left(\int_p A_i^a T_j^a dx_i \right),$$

where p are the intersection points of Γ and Σ , and $\int_p$ stands for the integration over Γ with the starting point at p . The expression $T_j^a T_j^a$ is proportional to the unit matrix with coefficient the quadratic Casimir $c_j = j(j+1)$ of $SU(2)$. This observation led [5] to the conclusion that $A_j = \sqrt{j(j+1)}$. We shall show instead that formula (1) holds true.

The area operator $(E_i^a(x)E_i^a(x))$ contains a product of two fields $E_i^a(x)$ at the same point and potentially needs a regularization. The answer $c_j = j(j+1)$ is certainly correct to the leading order in j which might be used as the parameter in the semi-classical expansion. However, there might be quantum corrections to this formula similar to the shift by $\hbar\omega/2$ in the energy spectrum of the harmonic oscillator $E_n = \hbar\omega(n + \frac{1}{2})$.

First, let us mention that the quantization problem for polynomials of commuting variables $[t^a, t^b] = 0$ which after quantization $t^a \mapsto T^a$ acquire Lie algebra commutation relations $[T^a, T^b] = f_{abc}T^c$ has a long history in mathematics. In fact, there is a universal *quantization map* $\mathcal{Q}$ which was discovered by Harish-Chandra for semi-simple Lie algebras and then generalized by Duflo [10] to the case of arbitrary Lie algebras. The main property of $\mathcal{Q}$ is that given two Casimir elements α and β (a particular example of a Casimir element is the quadratic Casimir $T^a T^a$) the product of quantizations $\mathcal{Q}(\alpha)\mathcal{Q}(\beta)$ coincides with the quantization of the product, $\mathcal{Q}(\alpha\beta)$. In more detail,

$$\mathcal{Q}(\alpha) = \text{Sym} \left[\det \left(\frac{\sin(x)}{x} \right)_{x=T_{\text{ad}}^a \frac{\partial}{\partial t^a}} \alpha(t) \right],$$

where T^a are generators in the adjoint representation, and Sym stands for the symmetric quantization map, *e.g.* $\text{Sym}(t^a t^b) = \frac{1}{2}(T^a T^b + T^b T^a)$. For instance, in the case of the Lie algebra $su(2)$, $\mathcal{Q} = \text{Sym} (1 + \frac{1}{12} \frac{\partial^2}{\partial t^a \partial t^a} + \dots)$, where $\dots$ stands for the terms containing higher derivatives in t^a . When applying $\mathcal{Q}$ to $c = t^a t^a$ one obtains $\mathcal{Q}(c) = T^a T^a + \frac{1}{4}$ with the eigenvalue $\mathcal{Q}(c)_j = (j + \frac{1}{2})^2$ in the spin j representation. Note that the minimal value of the Casimir corresponding to the (infinite-dimensional) representation of spin $j = -\frac{1}{2}$ is equal to zero.

Second, instead of using the A -representation one can rewrite the wave functional $W_\Gamma(A)$ in the E -representation by means of the functional Fourier transform,

$$\tilde{W}_\Gamma(E) = \int \mathcal{D}A e^{i \int \text{Tr}(EA)} W_\Gamma(A). \quad (4)$$

This expression can be simplified using the geometric quantization formula for the Wilson line [11,12],

$$W_\Gamma(A) = \int \mathcal{D}g e^{i \int_\Gamma \text{Tr}(\tau g^{-1} \partial_s g ds + A(g\tau g^{-1}))}, \quad (5)$$

where the auxiliary field g is a group valued function on the contour Γ , τ is a constant diagonal matrix, and s is a parameter along Γ . The right hand side of (5) is well-defined only when the eigenvalues of p are integers or half-integers [12]. Putting together the functional Fourier transform (4) and the geometric quantization (5) yields,

$$\tilde{W}_\Gamma(E) = \int dg \delta(E - g\tau g^{-1} \delta_\Gamma), \quad (6)$$

where δ_Γ stands for the δ -function supported on Γ . Equation (6) suggests that E vanishes outside Γ . On the contour, we obtain $\sqrt{\text{Tr} E^2} = \sqrt{\text{Tr} \tau^2} \delta_\Gamma$. Hence, the contribution to the area operator coming from the transversal

intersection of Γ and Σ is given by $A_j = \sqrt{\text{Tr}\tau^2}$. The relation between τ and j depends on the regularization of the functional integral over g . Using the regularization of [11] one obtains $A_j = j + \frac{1}{2}$ (the regularization of [12] yields $A_j = j$).

Third, the same problem of regularization of the $\text{Tr}E^2$ operator arises in 2-dimensional Yang-Mills theory as considered by Witten [13]. There, the value of the regularized quadratic Casimir has an influence on the results for the volumes of the moduli spaces of flat connections on Riemann surfaces. The same volumes can be computed by a different mathematically rigorous procedure (see *e.g.* [14]). Once again, the correct regularization of the Casimir element is $c_j = (j + \frac{1}{2})^2$ [13].

Finally, upon proper use of the gauge constraints, the problem reduces to geodesic motion of a particle on the group manifold, in our case $SU(2)$. The operator $\text{Tr} E^2$ becomes the Hamiltonian of the particle. Quantum mechanically, for motion on a curved manifold, it is known that in addition to the Laplacian (which would reproduce the standard Casimir) a term proportional to the scalar curvature of the manifold should be added, with the coefficient fixed to $\frac{1}{8}$ in order to reproduce the right conformal properties. In our case, this term is a constant and contributes the extra $\frac{1}{4}$. Alternatively, one can reduce the system even further to the dynamics of the eigenvalues of the group element. It is known that these behave as free fermions on a circle, and the extra contribution is seen to be the ground state Fermi energy of the system.

We conclude that all the evidence is in favour of the formula (1) for the spectrum A_j of the elementary objects. This implies the Bekenstein-Mukhanov type discrete spectrum of the black hole area, and the discrete spectrum of the black hole radiation.

We now turn to the black hole thermodynamics. We consider a quantum black hole as consisting of a large number of identical elementary components. Examples of such elementary components are the Wilson lines in the Ashtekar's gravity and D₀-branes in M-theory. Different viewpoints on the degree of distinguishability of these components will be discussed, and the entropy will be calculated in each case. Viewing the set of elementary components as the constituents of a grand canonical system, thermodynamics determines the entropy S of the black hole, to be given by

$$S = k(A\beta + \ln Z), \quad (7)$$

where k is the Boltzmann constant, A is the conserved black hole area, β is a temperature-like parameter dual to the area and Z is the partition function,

$$Z = \sum_A \Omega(A) e^{-\beta A} = \sum_{A,N} \Omega(A,N) e^{-\beta A}. \quad (8)$$

Here $\Omega(A)$ is the multiplicity of states of area A , and $\Omega(A,N)$ is the multiplicity of state of area A and N elementary components. The summation is over all possible areas and number of components.

To proceed, we will consider the elementary components as identical but *completely distinguishable* independent quantum systems. Then the partition function becomes

$$Z = \sum_N Z_1^N = \frac{1}{1 - Z_1}, \quad (9)$$

where $Z_1 = \sum_A \Omega(A,1) e^{-\beta A}$ is the partition function of a single component. The above expression for Z implies that β can never be less than the *Hagedorn temperature* parameter β_o , fixed by the relation $Z_1(\beta_o) = 1$.

From equation (8) the area is related to the partition function by formula,

$$A = -\frac{d \ln Z}{d\beta} = -\frac{1}{1 - Z_1} \cdot \frac{dZ_1}{d\beta}.$$

Restricting our considerations to macroscopic areas $A \gg 1$ only, this relation implies that $1 - Z_1 \sim \frac{1}{A}$, and thus $\beta \rightarrow \beta_o$ and $\ln Z$ grows only as $\ln A$. The dominant contribution to the entropy will therefore be given by $S = k\beta_o A$. The entropy is always proportional to the area. This is a completely general result, valid for any system consisting of distinguishable components.

To fix the proportionality constant, we restrict our considerations to the case of the area spectrum given by equation (1), *i.e.* $A_n = n$, $n = 1, 2, 3, \dots$, and the degeneracy function $\Omega(A,1) = \Omega(n,1) = g(n) = n$. Note that the states enumerated by the spin quantum number j , taking integer or half-integer values, are now enumerated by $n = 2j + 1$, a positive integer. We obtain $Z_1 = (2 \sinh \frac{\beta}{2})^{-2}$. The partition function diverges at the Hagedorn temperature $\beta = \beta_o = \ln \frac{3+\sqrt{5}}{2}$ and the area becomes macroscopic as $\beta \rightarrow \beta_o$. In that limit the entropy becomes

$$S = \frac{kA}{4\pi\gamma l_p^2} \ln \left(\frac{3+\sqrt{5}}{2} \right), \quad (10)$$

where we have acknowledged that the area is measured in units of $4\pi\gamma l_p^2$, where l_p is the Planck length and γ is the Immirzi parameter [15].

Equation (10) represents our final result for the entropy of a black hole. The validity of this derivation depends crucially on two important claims: that equation (1) gives the true area spectrum and that the area

constituents, i.e. the edges in the Ashtekar's gravity approach, are completely distinguishable. Attempts to justify the former claim have already been made. Let us now turn to the latter.

The elementary components of a black hole could *a priori* be either *identical*, *partially distinguishable* or *completely distinguishable*. Completely distinguishable components would mean that: (1) There is a difference between assigning two given spin values j_1 and j_2 to two different edges according to (edge 1, edge 2) = (j_1, j_2) and (edge 1, edge 2) = (j_2, j_1) . (2) Even if $j_1 = j_2$, there is a difference between assigning two different spin states m_1 and m_2 to the two different edges.

In calculating the entropy of equation (10), the edges were considered to be *completely distinguishable*, i.e. both claims (1) and (2) are imposed. Since the positions of different edges are determined by the *spin network*, which is formed by the way in which the edges are connected to each other and to the outside world, this viewpoint appears to be the one best describing the physical model, and as such, the one we should adopt (see also [16]).

In papers [7,8] a different view is adopted: The edges are *partially distinguishable*. Indeed, the multiplicity formula $\Omega(\{N_n\}) = \prod_n g(n)^{N_n}$ applies if claim (1) above is *not* adopted while (2) still applies. Evaluating (8) with this counting of states we obtain

$$Z = \sum_{\{N_n\}} \Omega(\{N_n\}) e^{-\beta A} = \prod_n \frac{1}{1 - g(n) e^{-\beta A(n)}} \quad (11)$$

If each edge does not have an exponentially increasing density of states (in which case it would be itself a macroscopic black hole) the only poles of the above expression are at $e^{\beta A(n)} = g(n)$. Calling $\bar{\beta}_o$ the lowest of these values of β , occuring for some n_o , we deduce that for macroscopic areas the model will exhibit Bose condensation at the spin $j_o = (n_o - 1)/2$ and the entropy will be $S = \bar{\beta}_o A$. For our area function (1) and the degeneracy function $g(n) = n$ this gives an entropy $S = \frac{kA\gamma_0}{4l_p^2}$ with $\gamma_0 = \frac{\ln 3}{3\pi}$. Bose condensation occurs at spin $j = 1$. For Ashtekar's choice of area function the result is instead $\gamma_0 = \frac{\ln 2}{\pi\sqrt{3}}$, and Bose condensation occurs at $j = \frac{1}{2}$.

Finally, if the edges are *identical* we adopt neither claim (1) nor (2) and have the equivalent of a Bose-Einstein gas. Then, it turns out that the entropy is related to the area by $S \propto A^t$, where the exponent satisfies $t < 1$ if each edge does not have an exponential density of states. For the choices of area spectrum (1) and de-

generacy function $g(n) = n$, the exponent acquires the value $t = \frac{2}{3}$. No Bose condensation occurs.

In conclusion, the area law $S = \beta_o A$ and the appearance of a Hagedorn temperature are the main results. These are extremely generic, requiring only *some* distinguishability of the elementary components. Since the Immirzi parameter is not fixed by the quantum theory [15], the different values of β_o are of somewhat secondary importance. There is, however, a crucial physical difference between our result and the result of [7,8]. While the latter suggest that Bose condensation occurs and area fluctuations are strongly suppressed, in our fully distinguishable case no Bose condensation occurs and the area within any solid angle of the black hole will exhibit fluctuations of order $\sqrt{A}$.

Acknowledgements. We thank J. Baez, V. Mukhanov and T. Strobl for useful discussions.

-
- [1] R. Gambini, J. Pullin, *Loops, Knots, Gauge Theories and Quantum Gravity* Cambridge University Press, Cambridge, 1996.
 - [2] A. Ashtekar, *Lectures on non-perturbative canonical gravity* Advanced Series in Astrophysics and Cosmology, Vol.6, World Scientific, Singapore, 1991.
 - [3] T. Banks *et al.* JHEP **9801** (1998) 008; H. Liu, A. A. Tseytlin JHEP **9801** (1998) 010.
 - [4] C. Rovelli and T. Thiemann, Phys.Rev. **D57** (1998) 1009-1014.
 - [5] C. Rovelli and L. Smolin, Nucl.Phys. **B442** (1995) 593-622; S. Frittelli, L. Lehner and C. Rovelli, Class.Quant.Grav. **13** (1996) 2921-2932.
 - [6] J. D. Bekenstein, V. F. Mukhanov, Phys.Lett. **B360** (1995) 7-12.
 - [7] A. Ashtekar *et al.* Phys.Rev.Lett. **80** (1998) 904-907.
 - [8] S. Chaudhuri and D. Minic, Phys. Lett. **B433** (1998) 301-306.
 - [9] J.C.Baez, *An Introduction to Spin Foam Models of BF Theory and Quantum Gravity* preprint gr-gc/9905087.
 - [10] M. Duff, Ann. Sci. École Norm. Sup. **10** (1977) 265-288.
 - [11] H.B. Nielsen and D. Rohrlich, Nucl. Phys. **B299** (1988) 471.
 - [12] A. Alekseev, L. Faddeev and S. Shatashvili, J. Geom. Phys. **5**, no. 3 (1989) 391.
 - [13] Edward Witten, J.Geom.Physics **9** (1992) 303-368.
 - [14] E. Meinrenken and C. Woodward, Progr. Math. **172** (1999) 271-295.
 - [15] C. Rovelli and T. Thiemann, Phys.Rev. **D57** (1998) 1009-1014.
 - [16] A. Strominger, Phys. Rev. Lett. **71** (1993) 3397-3400.